

Bleeding in Early Pregnancy

If you need this leaflet in a different language or accessible format please speak to a member of staff who can arrange it for you.

اگر به این بروشور به زبان دیگر یا در قالب دسترس پذیر نیاز دارید، لطفاً با یکی از کارکنان صحبت کنید تا آن را برای شما تهیه کند.

Jeśli niniejsza ulotka ma być dostępna w innym języku lub formacie, proszę skontaktować się z członkiem personelu, który ją dla Państwa przygotowuje.

Dacă aveți nevoie de această broșură într-o altă limbă sau într-un format accesibil, vă rog să discutați cu un membru al personalului să se ocupe de acest lucru pentru dumneavoastră

如果您需要本传单的其他语言版本或无障碍格式，请联系工作人员为您安排。

إذا احتجت إلى هذه النشرة بلغة أخرى، أو بتنسيق يسهل الوصول إليه، يرجى التحدث إلى أحد الموظفين لترتيب ذلك لك.

This leaflet has been created to provide you with information about what to do if you bleed during your pregnancy.

Bleeding is a common complication, occurring in 2 in 10 pregnancies, many of which go on to be normal and healthy pregnancies.

Bleeding can also be associated with miscarriage or an ectopic pregnancy (a pregnancy growing outside of your womb, affecting 1 in 100 pregnancies).

The Emergency Department has recommended you contact the Buchanan Suite for further advice.

Attendance at the Buchanan Suite is by appointment only.

Please call to make an appointment even if you have been referred by your midwife, GP, A&E Doctor. Buchanan Suite staff will offer further advice and may then arrange an appointment for you to attend for a blood test or ultrasound scan.

**Buchanan Suite opening hours:
Monday to Friday 9.00am – 12.30pm
Monday & Wednesday 1.30pm – 5.00pm
(excluding Bank Holidays)
0151 290 4356 – Triage
0151 430 1737 – Clinic Reception**


Until you have been assessed, a definite diagnosis cannot be reached, and ectopic pregnancy cannot be ruled out.



If you have any of the following symptoms, please call the Gynaecology Ward 3E for advice 24 hours, 7 days a week on 0151 430 1522.

OR attend the Emergency Department

Abdominal pain, shoulder tip pain, pain in back passage, fainting or dizzy spells, heavy vaginal bleeding (soaking a pad every 30 - 60 minutes, passing large clots or flooding), feeling generally unwell.





**Mersey and West Lancashire
Teaching Hospitals**
NHS Trust

Whiston Hospital
Warrington Road,
Prescot, Merseyside, L35 5DR
Telephone: 0151 426 1600

St Helens Hospital
Marshall Cross Road,
St Helens, Merseyside, WA9 3DA
Telephone: 01744 26633

www.MerseyWestLancs.nhs.uk