

Colposcopy Examination

This leaflet can be made available
in alternative languages / formats on request.

*如有需要，本传单可提供其他语言/版式
此單張的其他語言/格式版本可按要求提供*

Na żądanie ta ulotka może zostać udostępniona
w innych językach/formatkach.

Why do I need a colposcopy examination?

We have invited you to have a colposcopy because of the result of your cervical screening test (previously called a 'smear test'). This is usually for one of 4 reasons, which are:

- we found some abnormal cells in your cervix and an infection with human papillomavirus (HPV).
- you have an HPV infection which has not gone away.
- you have had several screening tests where we were unable to give you a result (it is likely there is nothing wrong, but a colposcopy can find out for sure).
- the nurse or doctor who carried out your screening test thought your cervix did not look as healthy as it should.

It is important to remember that most people who have a colposcopy do not have cervical cancer.

The colposcopy examination

A colposcopy is an examination to check whether there are abnormal cells on your cervix, and if so, how serious they are. This is a follow-up to your cervical screening test. Colposcopy usually takes place in an outpatient hospital clinic.

A specialist will take a close look at your cervix using a magnifying lens with a light (a colposcope). They may take a small tissue sample (a biopsy) to check any areas of your cervix which look unusual. If the colposcopy confirms there are abnormal cells on your cervix, you may need to have them removed to help prevent cervical cancer.

Having a colposcopy

Before your appointment

If you would like a female nurse or doctor to carry out your colposcopy, please call the clinic to request this when you receive your appointment: **01744 646301**

If you think you might have your period on the day of your appointment, you may want to call the clinic to arrange a different appointment. But you can come for your appointment during your period if you want to: **01744 646301**.

To make it easier to look at your cervix during your colposcopy, please do not have sex or use vaginal medications, lubricants or creams for at least 24 hours beforehand.

You should take a panty liner to your appointment as you may have a small amount of vaginal discharge after your colposcopy. If you have a small tissue sample taken (a biopsy), you may have some bleeding too.

You can bring a friend, partner or member of your family with you if you want.

What happens before the examination?

First the doctor or nurse will want to ask you some questions. These questions relate to your periods, the type of contraception you use, any operations or illnesses you have had in the past, and so on.

You will then be taken into a separate changing room and asked to undress from the waist downwards (though a full skirt need not be removed).

What happens during the examination?


You will be taken into the examination room and lie down on a special colposcopy couch/bed with your knees bent. You will be asked to place your legs onto some padded supports. You will have a paper sheet or towel to cover your stomach and hips. The examination takes about 10 to 20 minutes.

Just like at your screening test, the nurse or doctor will put a speculum into your vagina and open it gently. They will then use a colposcope to take a close look at your cervix. The colposcope does not go inside you, or even touch your skin. It stays about 30cm (12 inches) outside your vagina. The image of your cervix from the colposcope will sometimes be on a screen. This helps the nurse or doctor see your cervix more clearly.

They will dab different liquids onto your cervix. The liquids make any abnormal cells a different colour so that they can be seen more easily. If the nurse or doctor finds anything unusual they may take away a small tissue sample, a few millimetres across (a biopsy). A biopsy is about the size of a pinhead. You may feel a slight stinging, but it should not be painful.

The biopsy will be sent to the laboratory for further tests.

- We will send you a letter to your home address telling you of the biopsy result in about 4-6 weeks or sooner.
- The letter will also have information on whether you will need further assessment or treatment in the Colposcopy Clinic, including a contact number if you have any questions or need to rearrange your appointment.
- The examination can feel uncomfortable and some people may feel some pain. If it feels painful, tell the nurse or doctor and they will try to make it more comfortable for you.



Having a colposcopy examination

After your appointment

Most people feel well enough to go about their day-to-day activities straight away, but some may need to go home and rest for a while. You may have some brownish discharge from your vagina from the liquids that were used during your colposcopy.

For the next few days, you may have some light bleeding from your vagina, especially if you have had a biopsy. This is normal and usually stops after 3 to 5 days. It's best to avoid sex, using tampons, and any vaginal medications, lubricants or creams until the bleeding stops.

What will the examination show?

The nurse or doctor may be able to tell you what they have found straight away. If you have had a biopsy taken, it will need to be checked in the laboratory. If this happens, you will get your results by post about 4 weeks later.

A normal result

About 4 in 10 people who have a colposcopy will have a normal result. If you have a normal colposcopy result, this means that your cervix looks healthy and you have low risk of developing cervical cancer before your next screening test.

You can have a normal colposcopy result even if you had an abnormal result in your cervical screening test.

Abnormal cells confirmed

About 6 in 10 people will have abnormal cells found at colposcopy. The medical terms for abnormal cells are CIN ('cervical intraepithelial neoplasia') and CGIN (cervical glandular intraepithelial neoplasia). CIN and CGIN are not cancer, but can sometimes go on to develop into cancer.

Your colposcopy and biopsy results will show if you need to have the abnormal cells removed or whether they can be left alone for now. This will depend on whether your CIN is 'low grade' or 'high grade' (see below):

CIN 1 ('low grade')

You are unlikely to develop cervical cancer. Often the abnormal cells will go away on their own when your immune system gets rid of the HPV. This happens in most cases. We will normally invite you for another cervical screening test in 12 months to check whether you still have HPV.

CIN 2 or CIN 3 ('high grade')

You have a higher chance of developing cervical cancer than a woman with 'low grade' CIN. We will normally offer you treatment to remove the abnormal cells as this will lower your risk of developing cervical cancer.

Cervical cancer

Rarely, someone having a colposcopy will be found to have cervical cancer. If this happens to you, we will refer you for care and treatment from a team of specialists. Cancers diagnosed through screening are usually found at an earlier stage. People who have early stage cancers are more likely to survive than people with later stage cancers.

What if I need treatment?

If your examination shows that you need treatment you will be offered the option to have it carried out during your visit to the colposcopy clinic. People sometimes need to come back for another colposcopy to have treatment, but often it is possible to remove the abnormal cells during your first colposcopy. You will receive information about the treatment before your visit. If you have not received this information, contact the Colposcopy clinic and we will be pleased to send it to you.

There are several equally effective methods available to treat CIN. The aim of all methods of treatment is to destroy all the cells affected by CIN, with the least disruption to normal tissue. We usually use a local anaesthetic for the treatment, so you will be awake but will not feel any pain.

The choice of treatment will depend on your particular case, the preferred method of the doctor doing the colposcopy, and on the methods available at the clinic.

You can be treated for most abnormalities as an outpatient and so you will not need to stay in hospital.

If you are pregnant, we will be able to remove the abnormal cells after you give birth. You should talk to the nurse or doctor about when it is best for you to have this done.

Risks of treatment

Although it is an effective way of preventing cervical cancer, treatment has some risks.

There is a risk of infection from having abnormal cells removed. Signs of infection that you need to see your GP about are:

- heavy bleeding.
- bleeding that does not go away.
- vaginal discharge that smells.
- pain in your tummy that doesn't go away.
- Having abnormal cells removed may affect any future pregnancies you have.

Women who get pregnant after having abnormal cells removed are not at increased risk of having their baby early if they undergo standard treatment. However if more cervical tissue needs to be removed, women are slightly more likely to have their baby 1 to 2 months early. This may affect around 16% of women (16 in 100) who have had this more extensive treatment and then have a baby.*

**Castanon, A and others (2014). Risk of preterm delivery with increasing depth of excision for cervical intraepithelial neoplasia in England: nested case-control study. British Medical Journal; 349: g6223*

Not everyone who has abnormal cells removed would have gone on to develop cervical cancer. We offer treatment to everyone with serious abnormal cells because it is not possible to tell who will and who will not develop cervical cancer.

Cancer can start to develop between your regular screening tests. It is important to look out for anything that is unusual for you, especially:

- bleeding between your periods, during or after sex, or after the menopause.
- a change to vaginal discharge.

If you have any of these changes, please see your GP as soon as possible. Usually these symptoms will **not** mean you have cancer. But if you are found to have cancer, getting it diagnosed and treated early can mean you are more likely to survive.

What happens to tissue samples after Colposcopy?

Your tissue samples will be kept by the laboratory for at least 10 years. Your colposcopy results may be seen by staff who work elsewhere in the health service, so that they can make sure the service is as good as possible and to improve the skills of specialist staff.

What should I do after treatment?

- If treatment was given following colposcopy you may have a blood-stained discharge for two to four weeks.
- During this time, and when you have your period, you will need to use sanitary towels rather than tampons.
- It is also best to avoid heavy exercise and **not** to have sexual intercourse.
- These measures allow the cervix to heal as quickly as possible.
- Treatment for CIN will have little or no effect on your future fertility, nor on your risk of having a miscarriage.

Will I need to have checkups?

Yes - It is important to keep your appointments to make sure that your cervix is still healthy.

After treatment you will be invited to have a cervical screening test sooner than usual to check that the treatment was successful at your GP practice unless stated otherwise. The interval duration of cervical screening test will depend on the type of abnormality you were treated for (CIN or CGIN).

Sometimes it is necessary for the colposcopy team to keep you under their care instead of discharging you back to your GP/community for subsequent follow up. The doctor or nurse colposcopist will be able to explain fully reasons for this if it applies to you.

If everything is satisfactory after your treatment and follow-up screening tests, you are usually advised to have screening carried out at 3 or 5 years depending on your age.

It is hospital policy not to give results over the telephone

- Please note this treatment may not be carried out during a period.
- If you are experiencing abnormal or heavy bleeding after treatment please telephone our clinic for advice – 01744 646301.

- Whilst under the care of the colposcopy clinic please do not attend your GP surgery for any smears as all necessary smears will be done here in the colposcopy clinic
- More information and support

For more information, you can:


- speak with your GP or practice nurse
- visit www.nhs.uk/cervical
- visit www.gov.uk and search for 'cervical screening'
- call the Jo's Cervical Cancer Trust helpline on 0800 802 8000

If you have any urgent problems outside of normal clinic hours, please contact:

Ward 3E on 0151 430 1522

Important Notice

If you are unable to attend for your appointment please telephone the clinic in advance so that your appointment can be given to someone else



Whiston Hospital
Warrington Road,
Prescot, Merseyside, L35 5DR
Telephone: 0151 426 1600

Twitter and Facebook
@mwlInhs
www.merseywestlancs.nhs.uk