

Students

There may be students present during your consultation as part of their on-going training. Please let the staff know if you wish to be seen by a doctor only.

Whiston Hospital
Warrington Road,
Prescot, Merseyside, L35 5DR
Telephone: 0151 426 1600



/sthknhs



@sthk.nhs

www.sthk.nhs.uk



St Helens and Knowsley
Teaching Hospitals
NHS Trust

Head Injury Advice

This leaflet can be made available
in alternative languages / formats on request.

如有需要，本传单可提供其他语言/版式
此單張的其他語言/格式版本可按要求提供

Na żądanie ta ulotka może zostać udostępniona
w innych językach/formatach.

Author: Department of Surgery
Department: Surgical Care
Document Number: STHK0853
Version: 3
Review date: 01 / 09 / 2024

Your doctor or nurse practitioner has examined you and is happy for you to go home. **However, for the first 24 hours at home you should have a responsible adult with you, who has read this information sheet.**

Most head injuries do not lead to serious complications. However, if you experience any of the following symptoms, you should go immediately to your nearest emergency department:

- severe or worsening headache (a mild headache is normal after a head injury)
- vomiting (being sick)
- confusion or feeling unusually sleepy
- fits (collapsing or passing out suddenly)
- any change to your eyesight
- dizziness or lack of co-ordination
- weakness in one or both of your arms or legs
- clear fluid or blood coming out of your ears or nose
- new deafness in one or both ears
- inability to remember what has happened (amnesia)

What you should not worry about

It is normal after a head injury to experience the following symptoms over the next few days:

- Mild headache. It is safe to take a painkiller such as paracetamol. (Please see the instructions on the packet for advice about how much to take and how often).
- Feeling sick (without vomiting) or not feeling hungry. This can be helped by drinking clear fluids (such as water) and avoiding drinking alcohol.
- Difficulty concentrating.
- Feeling more tired, which might then make you irritable or anxious.

These symptoms should disappear over the next two weeks.

However, if they don't or you are concerned about how you are feeling, please see your GP.

General advice

If you follow this advice you should get better more quickly and may relieve some of your symptoms.

- Try to avoid stressful situations as these can make any symptoms worse.
- Do not return to your normal school, college or work activity until you feel you have completely recovered.
- You should not drive a motor vehicle or operate machinery for at least 24 hours after your accident.
- Do not work on a computer or play computer games until you have completely recovered, as spending time concentrating on images or text on a screen may make your symptoms worse.
- Your ability to tolerate alcohol is reduced after a head injury, so you should not drink alcohol until you are completely recovered.
- Do not play any contact sports for at least 3 weeks and without talking to your doctor first.

Long-term problems

Most people recover quickly from their accident and experience no long-term problems. However, please contact your GP if you:

- are still experiencing problems two weeks after your accident
- are concerned as to whether you are fit to drive a car or motorbike
- develop new problems after a few weeks or months.