Who can I contact if I have a problem when I get home?

If you experience any problems related to your surgery or admission once you have been discharged home. Please feel free to contact 4A or 4B ward for advice from the nurse in charge. They will assist you via the telephone, advise you return to your GP or ask you to make your way to the ED department at Whiston Hospital depending upon the nature of your concern.

4A Ward – 0151 430 1420 **4B Ward (Monday to Friday only)** – 0151 430 1440



Self Catheterisation in Men

Frequently Asked Questions

This leaflet can be made available in alternative languages / formats on request.

如有需要,本传单可提供其他语言/版式 此單張的其他語言/格式版本可按要求提供

Na żądanie ta ulotka może zostać udostępniona w innych językach/formatach.

Whiston Hospital Warrington Road, Prescot, Merseyside, L35 5DR Telephone: 0151 426 1600 St Helens Hospital Marshall Cross Road, St Helens, Merseyside, WA9 3DA Telephone: 01744 26633



Author/Dept. Surgical Care Department Document Number: STHK1063 Version: 004 Review Date: 31/07/2021

What is self-catheterisation?

Self-catheterisation means that you insert a small, flexible plastic tube into the bladder. The urine flows out through this tube which is then removed and washed. By emptying the bladder regularly in this way, you will prevent a build-up of stagnant urine and should feel more comfortable. You will also be keeping the kidneys and bladder healthy by preventing urinary infections. Some people find that they have less difficulty preventing themselves becoming wet using this technique.

What equipment do I need?

- A catheter supplied at first by the outpatient clinic and then available to order on prescription from your GP
- Baby wipes

How do I do it?

- Wash your hands thoroughly
- Remove your pants and tuck other clothing out of the way
- Roll back the foreskin and wash the end of the penis using baby wipes or soap and water
- Hold the penis straight out from the body and, without touching the catheter on anything else, slowly and gently insert it until urine flows
- If the catheter seems to stick just before it enters the bladder, wait a few seconds to allow the sphincter (valve) muscle to relax and then push it gently in a little further
- Withdraw the catheter slowly
- Get yourself in a comfortable position , you can chose from sitting or standing by the toilet
- When the flow stops move a little to make sure that the bladder is completely empty
- Dispose of the catheter properly by putting it in a disposal bag before putting it in the household rubbish
- Wash your hands again

Personal Information

Use the catheter times a day

Discard the catheter and use a new catheter for each catheterisation (catheters are single use and are no re-usable)

Catheter Information

Catheter recommended	1	 	
(male)			

Size F

Are there any other important points?

This publication provides input from specialists, the British Association of Urological Surgeons, the Department of Health and evidence-based sources as a supplement to any advice you may already have been given by your GP. Alternative treatments can be discussed in more detail with your urologist or Specialist Nurse.