

Flexible Sigmoidoscopy

This leaflet can be made available
in alternative languages / formats on request.

如有需要，本传单可提供其他语言/版式
此單張的其他語言/格式版本可按要求提供

Na żądanie ta ulotka może zostać udostępniona
w innych językach/formatach.

Whiston Hospital
Warrington Road,
Prescot, Merseyside, L35 5DR
Telephone: 0151 426 1600

St Helens Hospital
Marshall Cross Road,
St Helens, Merseyside, WA9 3DA
Telephone: 01744 26633

Introduction

You have been advised by your GP or hospital doctor to have an investigation known as a flexible sigmoidoscopy.

This procedure requires your formal consent and this booklet has been written to enable you to make an informed decision in relation to agreeing to the investigation.

If you are unable to keep your appointment, please notify the department as soon as possible we will be able to arrange another date and time for you.

Reasons for and Benefits of the Procedure

Why do I need to have a flexible sigmoidoscopy?

You have been advised to undergo this investigation of the left side of your large bowel (colon) to help find the cause for your symptoms, thereby facilitating treatment, and if necessary to decide on further investigations.

There are many reasons for this investigation including:

- bleeding from the back passage
- abdominal pain and diagnosing the extent of some inflammatory bowel disease.
- follow-up inspection of previous disease
- assessing the clinical importance of abnormalities found on an x-ray or scan

What are the alternatives?

A CT scan examination is an alternative investigation to flexible sigmoidoscopy. It has the disadvantage that samples of the bowel cannot be taken if an abnormality is found. If this is the case a subsequent endoscopic examination may be required.

	<p>Omit your lunchtime dose.</p> <p>Resume your normal insulin with your evening meal.</p>		
<p>You should resume taking your normal tablets the morning after the procedure. However, your blood glucose may be higher than usual for a day or so.</p> <p>If you have any problems or require further advice contact your diabetes specialist.</p> <p>If you are a St Helens resident, contact 01744 646200 - option 5 for emergency advice.</p> <p>available 0800-2200 weekdays and 0800-1600 weekends and bank holidays.</p>			
<p>Continue usual basal rates and start bolus once normal diet started</p>		<p>Continue usual basal rates and start to bolus once normal diet started</p>	
<p>Continue usual basal rates, if having hypoglycaemic episodes set temporary basal rate of 80%, bolus based on carbohydrate content of replacement fluids</p>		<p>Insulin Pump</p>	

How to adjust my insulin before a Flexible Sigmoidoscopy

Insulins	Day before the procedure Monitor your blood glucose 4 times a day or if you feel hypoglycaemic	Time of Procedure:	
		Morning appointment	Afternoon appointment
3,4, or 5 injections daily (eg. an injection of mixed insulin 3 times a day or 3 meal time injections of short acting insulin and once or twice daily background)	Reduce dose of short acting and mixed insulins by 50% If carb counting, use the carbohydrate content of replacement fluids and calculate dose Reduce long acting insulin dose by 20%	Omit your morning dose of short acting insulin if no breakfast is eaten. If you normally take a long acting basal insulin in the morning you should take 80% of your normal dose. If you normally take a pre-mixed insulin the dose should be halved.	Take usual morning insulin dose if breakfast is eaten. Omit lunchtime dose. Your blood glucose will be checked on admission. Resume your normal insulin with your evening meal.

What is a flexible sigmoidoscopy?

This test is a very accurate way of looking at the lining of the left side of your large bowel.

The instrument used in this investigation is a flexible tube which enables light to be directed onto the lining of your bowel and relay pictures back to the endoscopist onto a television screen.

During the investigation the endoscopist may need to take some samples (biopsies) from the lining of your colon for analysis.

These will be retained. Photographs can be taken for record and documentation purposes.

What about my medication?

Routine Medication

If you have not received specific advice regarding these medications then you should continue to take them as normal.

It would be helpful to bring a list of your current medications.

Iron supplements and anti-diarrhea medications

Iron tablets must be stopped one week prior to your appointment.

If you are taking stool bulking agents (e.g. Fybogel, Regulan, Procto-fibe), Loperamide (Imodium), Lomotil or Codeine Phosphate you must stop these three days prior to your appointment.

Diabetics

If you are diabetic controlled on insulin or medication, please ensure the Endoscopy Department is aware so that the appointment can be made at the beginning of the list. Please see the guidelines printed at the back of the book.

Anticoagulants and Anti-platelet therapy

Please telephone the unit (01744 646231 between 9-10am or 2-3pm) if you are taking any anticoagulants (blood thinning medication) e.g. warfarin, dabigatran, apixaban or rivaroxaban or edoxaban.

Please stop Clopidogrel / Ticagrelor for 7 days prior to this test. If you have had a coronary stent inserted in the past 12 months please contact the unit.

Preparing for the investigation

Oral bowel Cleansing Solutions ('bowel prep')

You have been prescribed an oral bowel cleansing solution (sometimes also called a 'bowel prep') to clear out your bowels.

This is important to ensure the safety and effectiveness of the planned procedure. Diarrhoea is an expected outcome of treatment. Please ensure that you have ready access to a toilet at all times following taking the oral bowel preparation before the effects wear off.

Some common side effects of oral bowel preparation include nausea, vomiting, bloating and abdominal pain. There is also a risk of developing dehydration, low blood pressure or kidney problems with this medication. The clinician prescribing the oral bowel cleansing solution will have assessed your risk and identified the most appropriate medication for you.

<p>Twice daily (eg. Novomix 30[®], Humulin M3[®], Insuman comb 15[®], Insuman comb 25[®], Insuman comb 50[®], Humalog Mix 25[®], Humalog Mix 50[®])</p>	<p>Reduce dose by 50% (for example if you are on 24 units in the morning and 16 units in the evening, take 12 units and 8units instead)</p>	<p>Halve your usual dose Your blood glucose will be checked on admission Resume your normal insulin with your evening meal</p>	<p>Halve your usual dose Your blood glucose will be checked on admission Resume your normal insulin with your evening meal</p>
<p>You should resume taking your normal tablets the morning after the procedure. However, your blood glucose may be higher than usual for a day or so.</p> <p>If you have any problems or require further advice contact your diabetes specialist.</p> <p>If you are a St Helens resident, contact 01744 646200 - option 5 for emergency advice. available 0800-2200 weekdays and 0800-1600 weekends and bank holidays.</p>			

Insulins	Day before the procedure Monitor your blood glucose 4 times a day or if you feel hypoglycaemic	Time of Procedure:	
		Morning appointment	Afternoon appointment
Once daily (evening) (eg. lantus®/Glargine or Levemir/Detemir® or Degludec/Tresiba® or Insulatard® or Humulin I®)	Your dose will need to be reduced by 20% (for example, if you are on 30 units, take 24 units instead)	No dose adjustment necessary	No dose adjustment necessary
Once daily (morning) (eg. antus®/Glargine or Levemir/Detemir® or Degludec/Tresiba® or Insulatard® or Humulin I®)	Your dose will need to be reduced by 20% (for example, if you are on 30 units, take 24 units instead)	No dose adjustment necessary	Your dose will need to be reduced by 20% and your blood glucose will be checked on admission

The prescribed dose of oral bowel cleansing solution should not be exceeded.

The oral bowel cleansing solution should not be taken over a period of longer than 24 hours.

Oral bowel cleansing solutions predispose to dehydration. You should maintain a good fluid intake whilst taking these medications, ideally in the form of water. If you develop symptoms of dehydration, and cannot increase your fluid intake, then you should seek medical attention. These symptoms include dizziness or light-headedness (particularly on standing up), thirst or a reduced urine production.

Some medicines may need to be stopped around the time of your procedure. Your doctor will review your medication at the time they refer you for an investigation and will inform you if you need to temporarily stop taking any of your usual medicines. The medications your doctor may consider stopping are in the following groups; NSAIDs (anti-inflammatory), ACE inhibitors (for heart or blood pressure problems) and diuretics ('water tablets').

If you have not received specific advice regarding your regular medications then you should continue to take them as normal.

Patients taking immunosuppression medication will be given advice from their doctor before taking any bowel prep. If you are taking the oral contraceptive pill you will need to use additional precautions in the following week as the absorption of the pill may be affected.

Eating and drinking

It is necessary to have clear views of the lower bowel.

Therefore for 2 days before your appointment you will need to be on a low fibre diet and considerably increase your fluid intake.

Low Fibre Diet

Fibre is the indigestible part of cereals, fruit and vegetables.

Please follow the instructions on when to stop eating and drinking during your bowel preparation. If you have been prescribed a bowel preparation other than Plenvu, please follow the alternative instructions given.

Foods allowed	Foods to be avoided
White meat, skinless chicken, grilled or poached fish	Red meat, sausages, black/ white pudding and pies
Cheese, eggs, tofu	Fruit / Salad / Vegetables
Plain ice-cream, custard, jelly (No red or blackcurrant jelly),	Potato skins or chips
Boiled sweets	Wholemeal or granary bread
Butter, margarine, shredless marmalade or seedless jam	Wholemeal pasta, brown rice
White bread, pasta, rice, boiled or mashed potatoes (No skins)	Nuts and pulses, including baked beans
Clear soups	Puddings containing fruit or nuts
Water, fizzy drinks, fruit squash (Not blackcurrant)	Cakes
Tea or coffee (without milk)	Biscuits
	Yoghurts
	Breakfast cereal

You should resume taking your normal tablets the morning after the procedure.

However, your blood glucose may be higher than usual for a day or so.

If you have any problems or require further advice contact your diabetes specialist.

If you are a St Helens resident, contact 01744 646200 - option 5 for emergency advice (available 0800-2200 weekdays and 0800-1600 weekends and bank holidays).

Tablets	Day before the procedure	Time of Procedure:	
		Morning Appointment	Afternoon Appointment
DPP-IV inhibitors (eg. sitagliptin, saxagliptin, vildagliptin, alogliptin, linagliptin)	Take at breakfast time	Take as normal after procedure	Restart after procedure
GLP-1 analogue (eg. exenatide, liraglutide, lixisenatide)	Omit medication	Take as normal after procedure	Omit
SGLT-2 inhibitors (eg. dapagliflozin, canagliflozin, empagliflozin)	Omit medication	Take as normal after procedure	Omit

How do I take the Plenvu?

- Each box of Plenvu contains three sachets marked DOSE 1 and DOSE 2 (Sachet A + B).
- Open the sachet labelled DOSE 1. Empty the contents into a container. Add 500ml of water (not chilled). Stir until dissolved. This may take up to 8 minutes.
- Drink the contents of the container over 30 minutes followed by 500ml of clear fluid over 30 minutes.
- Acceptable clear fluids are water, diluted cordials (not blackcurrant), herbal tea, black tea or coffee.
- When you are due to take DOSE 2 open sachets A + B. Empty the contents in to a container. Add 500ml of water and stir until dissolved. As above this can take up to 8 minutes.
- Drink the contents over 30 minutes followed by 500ml of clear fluid over 30 minutes.
- If you are taking an oral medication take it an hour either side of your dose of preparation.

When do I take my bowel preparation?

For a morning appointment:

On the day before your examination

Following breakfast, even if you feel hungry do not have anything to eat after 9am, because your bowel must be completely empty for the examination. You can have clear fluids e.g. water, soft drinks (without bits but not blackcurrant juice), sieved soup, or black coffee or tea (without milk). Add sugar if you like. You should take the whole dose of Plenvu the day before your examination.

Tablets	Day before the procedure	Time of Procedure:	
		Morning Appointment	Afternoon Appointment
Acarbose	Omit medication	Omit morning dose if you have been told to fast from midnight	Omit morning and lunch time doses and restart from next dose after procedure
Meglitinide (eg. repaglinide or nateglinide)	Omit medication	Omit morning dose if you have been told to fast from midnight	Take your morning dose if eating breakfast. Do not take your lunchtime
Metformin / Glucophage MR	Omit medication after breakfast dose	If taken once a day – take after procedure If taken twice a day – take first dose after the procedure	If taken once a day – take after procedure If taken twice a day – omit morning dose and take evening dose after procedure

7pm

Make up the first dose of 500ml of Plenvu and drink it over 1-2 hours. You will experience increasingly watery stools, this is normal, so ensure you can get to a toilet easily.

It is important to drink an additional 500ml of water or clear fluids during the evening.

Allow 2 hours for the Plenvu to work after finishing the first dose.

When you are no longer experiencing watery stools go to bed as usual.

The morning of your examination

Do not forget do not eat solid food but continue to drink clear fluids until 2 hours before your appointment.

6am

Make up the second dose of 500ml of Plenvu and drink it over 1-2 hours.

When you have finished this allow 2 hours for the Plenvu to work.

Again you will experience increasingly watery stools, this is normal.

It is important to drink an additional 500ml of water or clear fluids during the morning.

When you are no longer experiencing watery stools, start your journey to the hospital for your appointment.

What happens when I arrive?

You should expect to be in the department for approximately three hours. You will be met by a healthcare professional who will ask you a few questions, one of which concerns your arrangements for getting home. You will be able to ask further questions about the investigation.

You will have brief medical assessment where a qualified endoscopy nurse will ask you some questions regarding our medical health to confirm that you are fit to undergo the investigation.

If you are diabetic, your blood glucose level will also be recorded.

The nurse will ensure you understand the procedure and discuss any further concerns or questions you may have. If you have not already done so, and you are happy to proceed, you will be asked to sign your consent form.

If you are having sedation, the nurse will insert a small cannula (small plastic tube) in your arm or hand through which sedation will be administered later.

Sedation and Analgesia

Intravenous sedation and analgesia

A sedative drug is offered for this procedure. However most patients are comfortable with Entonox or even no medications.

Sedation will make you lightly drowsy and relaxed but not unconscious. You will be in a state called co-operative sedation.

This means that, although drowsy, you will still hear what is said to you and therefore will be able to follow simple instructions during the investigation.

One day before the procedure

If your procedure is scheduled for the **morning**:

- Following breakfast, do not eat any food after 9AM one day before the procedure.
- Take the bowel preparation as instructed. Usually two doses taken on the evening before the procedure.

If your procedure is scheduled for the **afternoon**:

- Following lunch, do not eat any food after 1PM one day before the procedure.
Take the bowel preparation as instructed. Usually one dose taken on the evening before the procedure and a second dose taken on the morning of the procedure.

Take additional clear fluid, and sugary drinks such as Lucozade or clear fruit juice to maintain the blood glucose levels.

On the day of the procedure

If your procedure is scheduled for the **morning**:

- You may drink clear fluids such as black tea or coffee, sugar free squash or water up to 2 hours before your procedure (ie 7AM).

If your procedure is scheduled for the **afternoon**:

- You may drink clear fluids such as black tea or coffee, sugar free squash or water up to 2 hours before your procedure (ie 11AM).

Important things to bring to hospital with you

- Glucose tablets or a sugary drink.
- Blood glucose testing equipment (if you usually monitor your blood glucose).

Hypoglycemia Advice

If you have any symptoms of a low blood sugar such as sweating, dizziness, blurred vision or shaking please test your blood sugar if you are able to do so.

If your blood sugar is less than 6mmol/L take 4 glucose tablets or 150ml of a sugary drink.

Please tell staff at the hospital that you have done this.

- Following your procedure you will be advised when it is safe to eat and drink.
- If you are eating and drinking normally you should resume taking your normal tablets.

Your blood glucose levels may be higher than usual for a day or so.

Eating and Drinking Whilst Preparing for a Colonoscopy

Two days before the procedure

- Follow the low residue diet instructions on page 6.
- Take your diabetes medications and/or insulin as normal.
- Test your blood glucose levels before administering insulin.
- Try to make sure you eat your usual amounts of carbohydrate from the low residue diet sheet.

Take additional clear fluid, and sugary drinks such as Lucozade or clear fruit juice to maintain the blood glucose levels.

If you have sedation you will not be permitted to drive home or use public transport alone, so you must arrange for a family member or friend to collect you. If you live alone, arrange for someone to stay with you, or if possible, arrange to stay with your family or a friend for about 24 hours.

Please note if you have had sedation you must not drive, take alcohol, operate heavy machinery or sign any legally binding documents for 24 hours following the procedure and you will need someone to accompany you home.

Older patients and those who have significant health problems (for example, people with significant breathing difficulties due to a bad chest) may be assessed by a doctor before being offered sedation.

Entonox

Entonox is a gas that you may be offered to help you manage your discomfort/pain during some bowel investigations.

It consists of 50% Oxygen and 50% Nitrous Oxide.

Nitrous Oxide is sometimes used for anesthetics but in this combination it works as a short acting painkiller.

Entonox is sometimes called 'gas and air' and is most commonly used during child birth.

As Entonox is short acting it can be used as a painkiller for any treatment that requires short - term pain relief.

Are there any side effects?

Entonox can make you feel drowsy and a little light-headed. Other possible side effects include dizziness, nausea and a dry mouth. All these sensations disappear rapidly after you stop using Entonox.

How do I use it?

As part of your admission, the nurse looking after you will complete a checklist to make sure it is appropriate for you to have Entonox. You will be shown the Entonox equipment and the nurse will explain how to use it.

Please tell the nurse if you have recently had:

- Eye surgery (with gas bubble insertion)
- Head injury with impaired consciousness
- Facial injuries
- Heavy sedation
- Previous laryngectomy
- Pneumothorax (collapsed lung), lung surgery, chest drain Severe Chronic Pulmonary Disease (breathing problems)
- Extensive swelling of the abdomen or bowel obstruction (blockage)
- Decompression sickness (recent dive)
- Air embolism
- Are pregnant or believe you may be pregnant

What can or cannot I do afterwards?

The effects should wear off quickly. However, we advise that you should be cautious about getting up and walking around until your balance has fully returned. You must wait a minimum of 30 minutes before attempting to drive or operate machinery.

Guidelines for people with diabetes undergoing Flexible Sigmoidoscopy

Instructions for your appointment

Please check your appointment information. Your appointment should be early morning. If not, please contact the endoscopy unit on 0151 430 1293 for an alternative appointment time.

How to adjust Diabetes Treatment

Whilst you are preparing for a flexible sigmoidoscopy your diet will be different and involve a period of fasting. This means your diabetes medications may need to be adjusted.

If you take medications to control Diabetes, please follow the instructions in the table on pages 20-23:

‘How to adjust my non-insulin diabetes medication before a flexible sigmoidoscopy ’

If you take insulin to control Diabetes, please follow the instructions in the table on page 24-27:

‘How to adjust my insulin before a flexible sigmoidoscopy ’

Please check the tables carefully. If you have any problems or require further advice contact your diabetes specialist.

If you are a St Helens resident, contact 01744 646200, select option 5 for emergency advice (available 0800-2200 weekdays and 0800-1600 weekends and bank holidays).

Before you leave the department, the endoscopist or nurse will explain the findings and any medication or further investigations required.

They will also inform you if you require further appointments.

You will be given an aftercare leaflet containing instructions if you have any problems following your procedure.

You can request a copy of the endoscopy report if you wish and a copy will also be sent to your GP.

If you have had sedation, the drug remains in your blood system for about 24 hours and you may feel drowsy later on, with intermittent lapses of memory.

If the person collecting you leaves the department, the nursing staff will telephone them when you are ready for discharge.

General points to remember

- If you are unable to keep your appointment please notify the endoscopy unit as soon as possible.
- It is our aim for you to be seen as soon as possible after your appointment time. However, there may be delays
- Emergencies can occur and these patients are given priority.
- The hospital cannot accept any responsibility for the loss or damage to personal property during your time on these premises.
- Please follow carefully the instructions on preparation, medications and pacemakers.

Frequently asked Questions (FAQs)

Hopefully this leaflet has answered all your questions.

Our website does have further information and includes a number of questions asked by other patients.

This is a legal requirement and you will not be insured by your motor insurance company if you do so.

Are there any benefits of using Entonox?

1. You will be in control.
2. The pain relief effects are rapid.
3. There are no long lasting side effects.

Are there any risks of using Entonox?

There is minimal risk involved with the administration of Entonox.

The main concern is that the gas might get trapped in the body: this is only a problem if you have specific medical conditions.

The flexible sigmoidoscopy investigation

In turn you will be escorted into the procedure room where the endoscopist and the nurses will introduce themselves and you will have the opportunity to ask any final questions.

The nurse looking after you will ask you to lie on your left side. They will then place the oxygen monitoring probe on your finger. The sedative/ analgesic drugs will be administered or you will be shown how to use the Entonox.

The flexible sigmoidoscopy involves maneuvering the endoscope around the left side of the bowel of your large bowel. There are some bends that naturally occur in the bowel and negotiating these may be uncomfortable for a short period of time but the sedation and analgesia will minimise any discomfort. Air is gently passed into the bowel during the investigation to facilitate the passage of the endoscope.

During the procedure, samples may be taken from the lining of your bowel for analysis.

Risks and discomforts of the Procedure

Lower gastrointestinal endoscopy is classified as an invasive investigation and because of that it has the possibility of complications. These occur extremely infrequently; we would wish to draw your attention to them.

The clinician who has requested the test will have considered this very carefully and the risks must be compared to the benefit of having the procedure carried out.

Frequent risks

- Abdominal discomfort (2 people in every 100)
- The procedure cannot be completed (1 person in every 10)

Serious risks

Uncommon (1 person in every 50-1000):

- Complications of over-sedation (problems with breathing, heart rate and blood pressure, which are short lived)
- Adverse reactions to drugs or equipment
- Bleeding from biopsy site (<1 person in every 1000) or polypectomy site (2 people in every 100), possibly requiring a blood transfusion
- Perforation (1 person in every 1000) or higher (3 people in every 1000) with polypectomy, requiring surgery including possible stoma.

Rare (1 person in every 1000-10,000):

- Cardiac irregularities including heart attack and stroke, requiring admission.

Very rare:

- Death (1 person in every 15,000)

What is a polyp?

A polyp is a protrusion from the lining of the bowel. Some polyps are attached to the intestinal wall by a stalk, and look like a mushroom, whereas others are flat without a stalk. Polyps when found are generally removed or sampled by the endoscopist as they may grow and later cause problems. Flat polyps are generally a little more difficult to remove.

Polypectomy

Polyps may be removed using a biopsy (cupped forceps) or a snare (wire loop). An electrical current known as diathermy may be used to cauterize the polyp as it is removed.

Smaller polyps may be removed using a biopsy or a snare.

For larger polyps a snare is placed around the polyp, diathermy is then applied and the polyp is removed.

Flat polyps (without any stalk) can be removed by a procedure called EMR (Endoscopic Mucosal Resection). This involves injecting the lining of the bowel that surrounds the flat polyp. This raises the area and allows the wire loop snare to capture the polyp.

Please let us know before your procedure if you have a pacemaker (PPM) or implanted defibrillator (ICd) and bring the any details about your device with you on the day of the test.

After the Procedure

You will be allowed to rest for as long as is necessary. Your blood pressure and heart rate will be recorded and if you are diabetic, your blood glucose will be monitored. Once you have recovered from the initial effects of the sedation/Entonox (which normally takes 30-60 minutes), you will be moved to a comfortable chair and offered a hot drink and biscuits.