

Exercises Following Transtibial (Below Knee) Amputation

This leaflet can be made available in alternative languages / formats on request.

如有需要,本传单可提供其他语言/版式 此單張的其他語言/格式版本可按要求提供

Na żądanie ta ulotka może zostać udostępniona w innych językach/formatach.

Author: Therapies

Department: Trauma & Orthopaedics

Document Number: STHK1095

Version: 001 Review Date: 31 / 08 / 2025

Introduction

This sheet has been designed to help you remember the exercises that you have been taught by your therapist. All of the exercises should be done slowly and smoothly. If you feel any pain, stop and tell your therapist or doctor.

Exercise helps to keep your strength and mobility as well as improve your blood flow.

These exercises will help to keep you independent either in a wheelchair or using an artificial leg (prosthesis).

Try to keep doing these exercises, even after you have been discharged from treatment.

This exercise sheet has been produced by P.I.R.P.A.G. (Physiotherapy Inter Regional Prosthetic Audit Group) Copyright © PIRPAG 2004

Static Quads

Push your legs straight out in front of you.

Push the back of your knees into the bed and tighten the thigh muscles.

Hold for 5 seconds.

Repeat this times.



Straight Leg Raise

With your legs out in front of you.

Tighten your thigh by squeezing the back of your knee into the bed.

Keeping you knee straight, lift your leg off the bed.

Hold for 5 seconds then slowly lower

Repeat ___ times

Repeat the above with the other leg.



Inner Range Quads

Sit with both legs straight out in front of you Place a rolled up towel, blanket/pillow under your knee.

Straighten your knee.

Hold for 5 seconds.

Repeat ___ times.

Repeat the above with the other leg.





Hip Adduction With Resistance

Sit with both legs out in front of you.

Place a pillow or rolled up towel between your knees.

Squeeze your legs together.

Hold for 5 seconds.

Repeat ___ times.

NB This exercise can also be performed when sitting in a wheelchair or at the edge of a bed.



Outer Range Quadriceps

Sit on a chair or edge of the bed.

Place your hands on your lap.

Straighten one knee

Hold for 5 seconds.

Now bend your knee.

Repeat ___ times.

Repeat the above with the other leg.





Static Gluteal Contractions

In sitting or lying, keep both legs close together.

Squeeze your buttocks as tightly as possible.

Hold for 5 seconds.

Repeat ___ times.



Hip Flexor Stretch

Lie on your back, preferably without a pillow.

Bend your knee towards your chest and hold with your hands.

Push your opposite leg down flat on to the bed.

Hold for 30 – 60 seconds, then relax.

Repeat ___ times.

Repeat the above with the other leg.



Bridging

Lie on your back with your arms at the side.

Place a couple of firm pillows or rolled up blankets under your thighs.

Pull in your stomach, tighten your buttocks and lift your bottom up off the bed.

Hold for 5 seconds.

Repeat ___ times.

To make this exercise more difficult, place your arms across your chest as shown in the picture.





Hip Flexion & Extension in Side Lying

Lie on your side and bend the bottom leg.

Lift your top leg slightly and bend your knee fully towards your chest.

Straighten your knee and push your leg backwards.

Repeat ___ times.

NB Try not to let your hips roll forwards or backwards.

Repeat the above with the other leg.





Hip Abduction in Side Lying

Lie on your side and bend the bottom leg.

Keep hips and top leg in line with your body.

Slowly lift your top leg up, keeping your knee straight and slowly lower.

Repeat __ times.

NB Try not to let your hips roll forwards or backwards.

Repeat the above with the other Leg.





Knee Flexion in Prone Lying

Lie on your stomach.

Keeping your thigh on the bed, bend your knee as far as possible.

Hold for 5 seconds then straighten your knee.

Repeat ___ times.

Repeat the above with the other leg.





Whiston Hospital Warrington Road, Prescot, Merseyside, L35 5DR Telephone: 0151 426 1600

St Helens Hospital Marshall Cross Road, St Helens, Merseyside, WA9 3DA Telephone: 01744 26633

